

HAIRER'S RECONSTRUCTION THEOREM WITHOUT REGULARITY STRUCTURES

FRANCESCO CARAVENNA
University of Milano-Bicocca (Italy)

JOINT WITH LORENZO ZAMBOTTI

ONE WORLD PROBABILITY SEMINAR ~ 21 JAN 2021

0. SUMMARY


- Stochastic analysis & the Sewing Lemma
- Hairer's Reconstruction Theorem
(elementary approach, optimal assumptions)
- Examples and applications

1. THE SEWING LEMMA

Fix two continuous but non smooth functions

$$X, Y: [0,1] \rightarrow \mathbb{R} \quad (\text{possibly random})$$

How to define the product $f(t) = "X(t) \dot{Y}(t)"$?

 Define its integral $I(t) = "\int_0^t X(s) \dot{Y}(s) ds"$

Key problem in stochastic analysis ($Y = \text{Brownian motion}$)

Robust definition of $I(\cdot)$?

Local description


$$\underbrace{\int_s^t X(u) \dot{Y}(u) du}_{\text{}} = \underbrace{X(s)(Y(t)-Y(s))}_{\text{}} + o(|t-s|) \quad (\text{if smooth})$$

$$I(t) - I(s) = \underbrace{A(s,t)}_{\text{"GERM"}} + o(|t-s|) \quad (1)$$

Sewing Lemma [Gubinelli 04] [Feyel, de La Pradelle 06]

Equation (1) has a solution (unique!) if, for some $\gamma > 0$,

$$|A(s,t) - A(s,u) - A(u,t)| \lesssim |t-s|^{1+\gamma} \quad (s \leq u \leq t)$$



$A(\cdot, \cdot)$ is called coherent germ

- Basic germ $A(s, t) = X(s) (Y(t) - Y(s))$

$$|A(s, t) - A(s, u) - A(u, t)| = |(X(u) - X(s))(Y(t) - Y(u))|$$

- Assume $X \in \mathcal{C}^\alpha$, $Y \in \mathcal{C}^\beta$: $|u - s|^\alpha |t - u|^\beta \lesssim |t - s|^{\alpha + \beta}$

coherent

[Young 1936] $I(t) = \int_0^t X(u) \dot{Y}(u) du$ canonical for $\alpha + \beta > 1$

- $Y = B = \text{Brownian motion}$ $\beta = \frac{1}{2} - \Rightarrow$ Young requires $\alpha > \frac{1}{2}$

This rules out most interesting cases!

- SDE driven by Brownian motion (σ smooth)

$$" \dot{X}(t) = \sigma(X(t)) \dot{B}(t) "$$

Expect $X \in C^\alpha$ with $\alpha = \beta = \frac{1}{2} - \Rightarrow \alpha + \beta < 1$

Basic germ $A(s,t) = X(s) (B(t) - B(s))$ not coherent

- Enriched germ coherent! (Rough paths...)

$$\tilde{A}(s,t) = X(s) (B(t) - B(s)) + \underbrace{X'(s)}_{\sigma'(X(s)) \sigma(X(s))} \underbrace{" \int_s^t (B(u) - B(s)) \dot{B}(u) du "}_{\text{Ito, Stratonovich, ...}}$$

2. DISTRIBUTIONS

$d > 1$ $\left\{ \begin{array}{ll} \text{Functions} & \rightsquigarrow \text{Distributions} \\ \text{Sewing Lemma} & \rightsquigarrow \text{Reconstruction Theorem} \end{array} \right.$

$$I(t) - I(s) = A(s, t) + o(|t-s|) \quad (1)$$

- Local approximation of $I(t) - I(s)$ is the germ $A(s, t)$
- Local approximation of $f(t) := \dot{I}(t)$ is $\bar{F}_s(t) := \frac{\partial}{\partial t} A(s, t)$

$$\langle f, \mathbb{1}_{[s, t)} \rangle = \langle \bar{F}_s, \mathbb{1}_{[s, t)} \rangle + o(|t-s|) \quad (1')$$

$\nwarrow$ distributions $\nearrow$

$D := \{ \varphi : \mathbb{R}^d \rightarrow \mathbb{R}, \varphi \in C_c^\infty \} = \text{test functions}$

$D' := \{ f : D \rightarrow \mathbb{R} \text{ linear \& continuous} \} = \text{distributions}$

$\forall K \subseteq \mathbb{R}^d \text{ compact} \quad \exists r = r_K \in \mathbb{N} \quad \forall \varphi \in D: \text{supp}(\varphi) \subseteq K$


$$|\langle f, \varphi \rangle| \lesssim \|\varphi\|_{C^r} := \max_{|k| \leq r} \|\partial^k \varphi\|_\infty$$

Definition. A GERM $(F_x)_{x \in \mathbb{R}^d}$ is a family of distributions.

Problem. Given a germ $(F_x)_{x \in \mathbb{R}^d}$, can we find a distribution $f \in D'$ "locally approximated by F_x " ?

What do we mean by "locally well approximated by F_x "?

Fix a test function $\varphi \in \mathcal{D}$ with $\int \varphi \neq 0$



Rescaled test function

$$\varphi_x^\lambda(z) := \frac{1}{\lambda^d} \varphi\left(\frac{z-x}{\lambda}\right)$$

We can require the same as (1'): $(\varphi = \mathbb{1}_{[0,1)})$

$$\langle f, \varphi_x^\lambda \rangle = \langle F_x, \varphi_x^\lambda \rangle + o(1) \quad \text{as } \lambda \downarrow 0$$

(unif. for x in compact sets)


We require slightly more: $o(1) \rightsquigarrow O(\lambda^\gamma)$

Problem. Given a germ $(F_x)_{x \in \mathbb{R}^d}$, find $f \in D'$ such that

$$|\langle f - F_x, \varphi_x^\lambda \rangle| \lesssim \lambda^\gamma \quad (2)$$

for some $\varphi \in D$ and $\gamma > 0$ (unif. $\lambda \in [0, 1]$, x in compact)

Uniqueness is for free. Existence requires **coherence!**



$$|A(s, t) - A(s, u) - A(u, t)| \lesssim (t-s)^{1+\gamma}$$

$\Downarrow$

$$|\langle F_x - F_y, \varphi_x^\lambda \rangle| \lesssim \lambda^{-1} (|y-x| + \lambda)^{1+\gamma}$$

3. COHERENCE

Definition. Fix $\gamma \in \mathbb{R}$. A germ $(F_x)_{x \in \mathbb{R}^d}$ is γ -coherent if there is one test function $\varphi \in \mathcal{D}$ with $\int \varphi \neq 0$ s.t.

$$\forall K \subseteq \mathbb{R}^d \text{ compact: } \begin{cases} |\langle F_x - F_y, \varphi_y^\lambda \rangle| \lesssim \lambda^\alpha (|x-y| + \lambda)^{\gamma-\alpha} \\ \text{unif. for } x, y \in K \text{ and } \lambda \in [0, 1] \end{cases}$$

for some $\alpha = \alpha_K \leq 0$ such that $\alpha \leq \gamma$.

If $\alpha_K \equiv \alpha$ ($\forall K$) then $(F_x)_{x \in \mathbb{R}^d}$ is called (α, γ) -coherent.

Coherence: F_x and F_y are "close" when x and y are close

Example If an "old" germ $(A(s,t))_{s,t \in [0,1]}$ is γ -coherent

$$(|A(s,t) - A(s,u) - A(u,t)| \lesssim (t-s)^{1+\gamma})$$

the "new" germ $(F_s := \partial_t A(s,t))_{s \in [0,1]}$ is $(-1, \gamma)$ -coherent

Example Fix a function $f \in \mathcal{C}^\gamma$ with $\gamma \in (0, \infty)$.

The (degree $\lfloor \gamma \rfloor$)-Taylor polynomial of f is $(0, \gamma)$ -coherent

$$F_x(\cdot) = \sum_{|k| < \gamma} \partial^k f(x) \frac{(\cdot - x)^k}{k!} \quad |F_x(z) - F_y(z)| \lesssim (|x-y| + \underbrace{|z-y|}_{\lambda})^\gamma$$

4. THE RECONSTRUCTION THEOREM

Theorem. Let $F = (F_x)_{x \in \mathbb{R}^d}$ be a γ -coherent ($\gamma \in \mathbb{R}$).

Then (2) has a solution: there is $f \in D'$ s.t.

$\forall K \subseteq \mathbb{R}^d$ compact, $\forall \psi \in D(B(0,1))$:

$$|\langle f - F_x, \psi_x^\gamma \rangle| \leq C \cdot \begin{cases} \lambda^\gamma & \text{if } \gamma \neq 0 \\ (1 + |\log \lambda|) & \text{if } \gamma = 0 \end{cases} \quad \begin{matrix} \text{unif. for} \\ x \in K, \\ \lambda \in (0, 1] \end{matrix}$$

where $C = C(K, F, \gamma) \lesssim \|\psi\|_{C^2}$ for a suitable γ (*)

If $\gamma > 0$, then $f = R F$ is unique and linear.

- Coherence is necessary for the Reconstruction Theorem

- For $\delta \leq 0$ no uniqueness, linearity ok

(For $\delta = 0$ the bound $|\log \lambda|$ is optimal)

Coherent germs form a vector space ($\varphi \rightsquigarrow \text{any } \chi$)

- Local homogeneity of a germ: for suitable $\beta = \beta_K$

$$|\langle F_x, \varphi_x^\lambda \rangle| \lesssim \lambda^\beta \quad \text{unif. for } x \in K, \lambda \in (0, 1]$$

- τ in (*) satisfies $\tau > \max \{ -\alpha_{\overline{K}_2}, -\beta_{\overline{K}_2} \}$

5. REGULARITY STRUCTURES

Consider a singular stochastic PDE on $\mathbb{R}^d$ ($\rightsquigarrow$ Lorenzo)

In **RS** the (distributional) **solution** $h(t,x)$ of the PDE is **represented by a coherent germ** $H = (H_{t,x})$

$(\Pi_{t,x})_{t \geq 0, x \in \mathbb{R}^d}$ "model", $h \in \mathcal{D}^\gamma$ "modelled distribution"

$\rightsquigarrow$ γ -coherent germ $H = (H_{t,x} := \Pi_{t,x} h(t,x))_{t \geq 0, x \in \mathbb{R}^d}$

If $\gamma > 0$, the solution is **$h(t,x) = \mathcal{R} H$** (reconstruction of H)

The germs arising in **RS** have a graded structure

$$F_x(\cdot) = \sum_{a \in A} c^a(x) \pi_x^a(\cdot)$$

$\downarrow$
finite set

$\downarrow$
local "basis" of distributions

This structure allows to define e.g. "products"

Another key ingredient of **RS** is Schauder estimates

Coherence in **RS**:

$$|\langle F_x - F_y, \varphi_y^\lambda \rangle| \lesssim \sum_{a \in A, a < \gamma} \lambda^a |x - y|^{\gamma - a} \lesssim \lambda^\alpha (|x - y| + \lambda)^{\gamma - \alpha}$$

$\alpha := \min A$

Thanks!